

CBSE Class-10 Mathematics

NCERT solution

Chapter - 8

Introduction to Trigonometry - Exercise 8.2

1. Evaluate:

(i) $\sin 60^\circ \cos 30^\circ + \sin 30^\circ \cos 60^\circ$

(ii) $2 \tan^2 45^\circ + \cos^2 30^\circ - \sin^2 60^\circ$

(iii) $\frac{\cos 45^\circ}{\sec 30^\circ + \csc 30^\circ}$

(iv) $\frac{\sin 30^\circ + \tan 45^\circ - \csc 60^\circ}{\sec 30^\circ + \cos 60^\circ + \cot 45^\circ}$

(v) $\frac{5 \cos^2 60^\circ + 4 \sec^2 30^\circ - \tan^2 45^\circ}{\sin^2 30^\circ + \cos^2 30^\circ}$

Ans. (i) $\sin 60^\circ \cos 30^\circ + \sin 30^\circ \cos 60^\circ$

$$= \frac{\sqrt{3}}{2} \times \frac{\sqrt{3}}{2} + \frac{1}{2} \times \frac{1}{2}$$

$$= \frac{3}{4} + \frac{1}{4} = \frac{4}{4} = 1$$

(ii) $2 \tan^2 45^\circ + \cos^2 30^\circ - \sin^2 60^\circ$

$$= 2(1)^2 + \left(\frac{\sqrt{3}}{2}\right)^2 - \left(\frac{\sqrt{3}}{2}\right)^2$$

$$= 2 + \frac{3}{4} - \frac{3}{4} = 2$$

$$\begin{aligned}
 \text{(iii)} \quad & \frac{\cos 45^\circ}{\sec 30^\circ + \csc 30^\circ} \\
 &= \frac{\frac{1}{\sqrt{2}}}{\frac{2}{\sqrt{3}} + 2} = \frac{\frac{1}{\sqrt{2}}}{\frac{2+2\sqrt{3}}{\sqrt{3}}} \\
 &= \frac{1}{\sqrt{2}} \times \frac{\sqrt{3}}{2+2\sqrt{3}} = \frac{\sqrt{3}}{\sqrt{2} \times 2(\sqrt{3}+1)} \\
 &= \frac{\sqrt{3}}{\sqrt{2} \times 2(\sqrt{3}+1)} \times \frac{\sqrt{3}-1}{\sqrt{3}-1} \\
 &= \frac{\sqrt{3}(\sqrt{3}-1)}{\sqrt{2} \times 2(3-1)} \quad [\text{Since } (a+b)(a-b) = a^2 - b^2] \\
 &= \frac{\sqrt{3}(\sqrt{3}-1)}{4\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} \\
 &= \frac{3\sqrt{2} - \sqrt{6}}{8}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv)} \quad & \frac{\sin 30^\circ + \tan 45^\circ - \csc 60^\circ}{\sec 30^\circ + \cos 60^\circ + \cot 45^\circ} \\
 &= \frac{\frac{1}{2} + 1 - \frac{2}{\sqrt{3}}}{\frac{2}{\sqrt{3}} + \frac{1}{2} + 1} = \frac{\frac{\sqrt{3} + 2\sqrt{3} - 4}{2\sqrt{3}}}{\frac{4 + \sqrt{3} + 2\sqrt{3}}{2\sqrt{3}}} = \frac{3\sqrt{3} - 4}{3\sqrt{3} + 4} \\
 &= \frac{3\sqrt{3} - 4}{3\sqrt{3} + 4} \times \frac{3\sqrt{3} - 4}{3\sqrt{3} - 4}
 \end{aligned}$$

$$= \frac{27+16-24\sqrt{3}}{27-16} \quad [\text{Since } (a+b)(a-b) = a^2 - b^2]$$

$$= \frac{43-24\sqrt{3}}{11}$$

$$(v) \frac{5 \cos^2 60^\circ + 4 \sec^2 30^\circ - \tan^2 45^\circ}{\sin^2 30^\circ + \cos^2 30^\circ}$$

$$= \frac{5\left(\frac{1}{2}\right)^2 + 4\left(\frac{2}{\sqrt{3}}\right)^2 - (1)^2}{\left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2}$$

$$= \frac{5 \times \frac{1}{4} + 4 \times \frac{4}{3} - 1}{\frac{1}{4} + \frac{3}{4}} = \frac{\frac{5}{4} + \frac{16}{3} - 1}{\frac{4}{4}}$$

$$= \frac{15+64-12}{12} = \frac{67}{12}$$

2. Choose the correct option and justify:

$$(i) \frac{2 \tan 30^\circ}{1 + \tan^2 30^\circ} =$$

(A) $\sin 60^\circ$

(B) $\cos 60^\circ$

(C) $\tan 60^\circ$

(D) $\sin 30^\circ$

(ii) $\frac{1 - \tan^2 45^\circ}{1 + \tan^2 45^\circ} =$

(A) $\tan 90^\circ$

(B) 1

(C) $\sin 45^\circ$

(D) 0

(iii) $\sin 2A = 2 \sin A$ is true when A =

(A) 0°

(B) 30°

(C) 45°

(D) 60°

(iv) $\frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ} =$

(A) $\cos 60^\circ$

(B) $\sin 60^\circ$

(C) $\tan 60^\circ$

(D) None of these

Ans. (i) (A) $\frac{2 \tan 30^\circ}{1 + \tan^2 30^\circ}$

$$= \frac{2 \times \frac{1}{\sqrt{3}}}{1 + \left(\frac{1}{\sqrt{3}}\right)^2}$$

$$= \frac{2}{\sqrt{3}} \times \frac{3}{3+1} = \frac{\sqrt{3}}{2} = \sin 60^\circ$$

$$(ii) (D) \frac{1 - \tan^2 45^\circ}{1 + \tan^2 45^\circ} = \frac{1-1}{1+1} = \frac{0}{2} = 0$$

(iii) (A) Since $A = 0$, then

$$\sin 2A = \sin 0^\circ = 0 \text{ and } 2 \sin A = 2 \sin 0^\circ$$

$$= 2 \times 0 = 0$$

$$\therefore \sin 2A = \sin A \text{ when } A = 0$$

$$(iv) (C) \frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ}$$

$$= \frac{2 \times \frac{1}{\sqrt{3}}}{1 - \left(\frac{1}{\sqrt{3}}\right)^2} = \frac{\frac{2}{\sqrt{3}}}{1 - \frac{1}{3}}$$

$$= \frac{2}{\sqrt{3}} \times \frac{3}{3-1} = \sqrt{3} = \tan 60^\circ$$

3. If $\tan(A+B) = \sqrt{3}$ and $\tan(A-B) = \frac{1}{\sqrt{3}}$; $0^\circ < A+B \leq 90^\circ$; $A > B$, find A and B.

$$\text{Ans. } \tan(A+B) = \sqrt{3}$$

$$\Rightarrow \tan(A+B) = \tan 60^\circ$$

$$\Rightarrow A+B = 60^\circ \dots\dots\dots(i)$$

$$\text{Also, } \tan(A-B) = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \tan(A - B) = \tan 30^\circ$$

$$\Rightarrow A - B = 30^\circ \dots\dots\dots(ii)$$

On adding eq. (i) and (ii), we get,

$$2A = 90^\circ \Rightarrow A = 45^\circ$$

On Subtracting eq. (i) and eq. (ii), we get

$$2B = 30^\circ \Rightarrow B = 15^\circ$$

4. State whether the following are true or false. Justify your answer.

(i) $\sin(A + B) = \sin A + \sin B$

(ii) The value of $\sin \theta$ increases as θ increases.

(iii) The value of $\cos \theta$ increases as θ increases.

(iv) $\sin \theta = \cos \theta$ for all values of θ .

(v) $\cot A$ is not defined for $A = 0^\circ$.

Ans. (i) False, because, let $A = 60^\circ$ and $B = 30^\circ$

Then, $\sin(A + B) = \sin(60^\circ + 30^\circ) = \sin 90^\circ = 1$

And $\sin A + \sin B = \sin 60^\circ + \sin 30^\circ = \frac{\sqrt{3}}{2} + \frac{1}{2} = \frac{\sqrt{3} + 1}{2}$

$\therefore \sin(A + B) \neq \sin A + \sin B$

(ii) True, because it is clear from the table below:

θ	0°	30°	45°	60°	90°
$\sin \theta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1

Therefore, it is clear, the value of $\sin \theta$ increases as θ increases.

(iii) False, because

θ	0°	30°	45°	60°	90°
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0

It is clear, the value of $\cos \theta$ decreases as θ increases

(iv) False as it is only true for $\theta = 45^\circ$.

$$\Rightarrow \sin 45^\circ = \frac{1}{\sqrt{2}} = \cos 45^\circ$$

(v) True, because $\tan 0^\circ = 0$ and $\cot 0^\circ = \frac{1}{\tan 0^\circ}$

$$= \frac{1}{0} \text{ i.e. undefined.}$$