

CBSE Class 11 Biology
NCERT Solutions
Chapter 13
Photosynthesis in higher plants

1. By looking at a plant externally can you tell whether a plant is C_3 or C_4 ? Why and how?

Ans. One cannot distinguish whether a plant is C_3 or C_4 by observing its leaves and other morphological features externally. Unlike C_3 plants, the leaves of C_4 plants have a special anatomy called Kranz anatomy and this difference can only be observed at the cellular level. For example, although wheat and maize are grasses, wheat is a C_3 plant, while maize is a C_4 plant.

2. By looking at which internal structure of a plant can you tell whether a plant is C_3 or C_4 ? Explain.

Ans. Yes, the leaves of C_4 plants have a special anatomy called Kranz anatomy. This makes them different from C_3 plants. Special cells, known as bundle-sheath cells, surround the vascular bundles. These cells have a large number of chloroplasts. They are thick-walled and have no intercellular spaces. They are also impervious to gaseous exchange. All these anatomical features help prevent photorespiration in C_4 plants, thereby increasing their ability to photosynthesise.

3. Even though a very few cells in a C_4 plant carry out the biosynthetic – Calvin pathway, yet they are highly productive. Can you discuss why?

Ans. The productivity of a plant is measured by the rate at which it photosynthesises. The amount of carbon dioxide present in a plant is directly proportional to the rate of photosynthesis. C_4 plants have a mechanism for increasing the concentration of carbon dioxide. In C_4 plants, the Calvin cycle occurs in the bundle-sheath cells. The C_4 compound (malic acid) from the mesophyll cells is broken down in the bundle-sheath cells. As a result, CO_2 is released. The increase in CO_2 ensures that the enzyme RuBisCo does not act as an

oxygenase, but as a carboxylase. This prevents photorespiration and increases the rate of photosynthesis. Thus, C_4 plants are highly productive.

4. RuBisCo is an enzyme that acts both as a carboxylase and oxygenase. Why do you think RuBisCo carries out more carboxylation in C_4 plants?

Ans. The enzyme RuBisCo is absent from the mesophyll cells of C_4 plants. It is present in the bundle-sheath cells surrounding the vascular bundles. In C_4 plants, the Calvin cycle occurs in the bundle-sheath cells. The primary CO_2 acceptor in the mesophyll cells is phosphoenol pyruvate – a three-carbon compound. It is converted into the four-carbon compound oxaloacetic acid (OAA). OAA is further converted into malic acid. Malic acid is transported to the bundle-sheath cells, where it undergoes decarboxylation and CO_2 fixation occurs by the Calvin cycle. This prevents the enzyme RuBisCo from acting as an oxygenase.

5. Suppose there were plants that had a high concentration of Chlorophyll-b, but lacked chlorophyll-a, would it carry out photosynthesis? Then why do plants have chlorophyll-b and other accessory pigments?

Ans. Chlorophyll-a molecules act as antenna molecules. They get excited by absorbing light and emit electrons during cyclic and non-cyclic photophosphorylations. They form the reaction centres for both photosystems I and II. Chlorophyll-b and other photosynthetic pigments such as carotenoids and xanthophylls act as accessory pigments. Their role is to absorb energy and transfer it to chlorophyll-a. Carotenoids and xanthophylls also protect the chlorophyll molecule from photo-oxidation. Therefore, chlorophyll-a is essential for photosynthesis.

If any plant were to lack chlorophyll-a and contain a high concentration of chlorophyll-b, then this plant would not undergo photosynthesis.

6. Why is the colour of a leaf kept in the dark frequently yellow, or pale green? Which pigment do you think is more stable?

Ans. Since leaves require light to perform photosynthesis, the colour of a leaf kept in the dark changes from a darker to a lighter shade of green. Sometimes, it also turns yellow. The production of the chlorophyll pigment essential for photosynthesis is directly proportional to the amount of light available. In the absence of light, the production of chlorophyll-a

molecules stops and they get broken slowly. This changes the colour of the leaf gradually to light green. During this process, the xanthophyll and carotenoid pigments become predominant, causing the leaf to become yellow. These pigments are more stable as light is not essential for their production. They are always present in plants.


7. Look at leaves of the same plant on the shady side and compare it with the leaves on the sunny side. Or, compare the potted plants kept in the sunlight with those in the shade. Which of them has leaves that are darker green? Why?

Ans. Light is a limiting factor for photosynthesis. Leaves get lesser light for photosynthesis when they are in shade. Therefore, the leaves or plants in shade perform lesser photosynthesis as compared to the leaves or plants kept in sunlight.


In order to increase the rate of photosynthesis, the leaves present in shade have more chlorophyll pigments. This increase in chlorophyll content increases the amount of light absorbed by the leaves, which in turn increases the rate of photosynthesis. Therefore, the leaves or plants in shade are greener than the leaves or plants kept in the sun.

8. Figure shows the effect of light on the rate of photosynthesis. Based on the graph, answer the following questions:

- (a) At which point/s (A, B or C) in the curve is light a limiting factor?**
- (b) What could be the limiting factor/s in region A?**
- (c) What do C and D represent on the curve?**



Ans.



(a) Generally, light is not a limiting factor. It becomes a limiting factor for plants growing in shade or under tree canopies. In the given graph, light is a limiting factor at the point where photosynthesis is the minimum. The least value for photosynthesis is in region A. Hence, light is a limiting factor in this region.

(b) Light is a limiting factor in region A. Water, temperature, and the concentration of carbon dioxide could also be limiting factors in this region.

(c) Point D represents the optimum point and gives the light intensity at which the maximum photosynthesis is recorded. The rate of photosynthesis remains constant after this point, even though the intensity of light.

9. Give comparison between the following:

(a) C_3 and C_4 pathways

(b) Cyclic and non-cyclic photophosphorylation

(c) Anatomy of leaf in C_3 and C_4 plants

Ans. (a) C_3 and C_4 pathways:

C_3 Pathway	C_4 Pathway
<p>i. In the C_3 pathway, the first product of carbon fixation is a 3 carbon atom compound 3-phosphoglyceric acid.</p> <p>ii. Photorespiration occurs in C_3 plants.</p>	<p>i. In the C_4 pathway, the first product of carbon fixation is oxaloacetic acid, a 4 carbon atom compound.</p> <p>ii. Photorespiration rarely occurs.</p>

iii. Plants are adapted to all climates except saline conditions.	iii. Plants are adapted to tropical climate and can tolerate halophytic conditions.
iv. This pathway is less expensive and requires only 18 molecules of ATP for the synthesis of one molecule of glucose.	iv. This pathway is more energy expensive and requires 30 molecules of ATP for the synthesis of one molecule of glucose.

(b) Cyclic and non-cyclic photophosphorylation:

Cyclic photophosphorylation	Non-cyclic photophosphorylation
i. It involves the participation of only PS I. ii. This system is not concerned with photo-oxidation of water. iii. It is only concerned with the production of ATP. iv. It occurs under special conditions when NADPH starts accumulating.	i. It involves the participation of both PS I and PS II. ii. The first step of this system is photo-oxidation of water resulting into H^+ , e^- and release of O_2 . iii. It is concerned with the reduction of $NADP^+$ and production of ATP. iv. It occurs under normal conditions.

(c) Anatomy of leaf in C_3 and C_4 plants:

Anatomy of leaf in C_3 plants	Anatomy of leaf in C_4 plants
i. Plant do not posses 'Kranz' anatomy of leaves. ii. Chloroplasts are not dimorphic. iii. Mesophyll cells have intercellular intercellular spaces.	i. Leaves of C_4 plants have 'Kranz' anatomy. ii. Chloroplasts are dimorphic. The chloroplasts of bundle sheath cells are larger in size and arranged centripetally. iii. Mesophyll cells do not have intercellular spaces.